

Forefoot and Midfoot ICD-10 Coding

Jeffrey D. Lehrman, DPM, FASPS, MAPWCA, CPC

Advisor, APMA Coding Committee

Advisor, APMA MACRA Task Force

Expert Panelist, Codingline

Fellow, American Academy of Podiatric Practice Management

Board of Directors, American Society of Podiatric Surgeons

Board of Directors, American Professional Wound Care Association

Editorial Advisory Board, *WOUNDS*

Twitter: [@DrLehrman](https://twitter.com/DrLehrman)

Hammertoe

- **M20.40** Other hammertoe(s) (acquired), unspecified foot
- **M20.41** Other hammertoe(s) (acquired), right foot
- **M20.42** Other hammertoe(s) (acquired), left foot
- **Q66.89** Other specified congenital deformities of feet

Acquired versus Congenital

Acquired


- Develop over a course of one's life.

Congenital

- Existing at birth or before birth.
- Develop during the first month of life

Hammertoe

- **M20.40** Other hammertoe(s) (acquired), unspecified foot
- **M20.41** Other hammertoe(s) (acquired), right foot
- **M20.42** Other hammertoe(s) (acquired), left foot
- **Q66.89** Other specified congenital deformities of feet



M20.41 Other hammertoe(s) (acquired)
right foot

M20.42 Other hammertoe(s) (acquired),
left foot

Q66.89 Other specified congenital
deformities of feet



M20.5X1 Other deformities of toe(s)
(acquired), right foot

M20.5X2 Other deformities of toe(s)
(acquired), left foot

Q66.89 Other specified congenital
deformities of feet



M20.5X1 Other deformities of toe(s)
(acquired), right foot

M20.5X2 Other deformities of toe(s)
(acquired), left foot

Q66.89 Other specified congenital
deformities of feet

Hallux Rigidus

–**M20.21** Hallux rigidus, right foot

–**M20.22** Hallux rigidus, left foot

Hallux Varus

–**M20.31** Hallux varus (acquired), right foot

–**M20.32** Hallux varus (acquired), left foot

Lisfranc Fracture Dislocation



Lisfranc Fracture Dislocation

- No one code
- So what do we have:
 - Displaced Fracture 2nd metatarsal
 - Dislocation of tarsometatarsal joint



- **Displaced fracture 2nd metatarsal**
 - S92.321__** Displaced fracture of second metatarsal bone, right foot
 - S92.322__** Displaced fracture of second metatarsal bone, left foot
- **Dislocation of tarsometatarsal joint**
 - S93.324__** Dislocation of tarsometatarsal joint of right foot
 - S93.325__** Dislocation of tarsometatarsal joint of left foot

**** All need 7th character**

7th character options

7th character options

Displaced fracture 2nd metatarsal

- A initial encounter for closed fracture
- B initial encounter for open fracture
- D subsequent encounter for fracture with routine healing
- G subsequent encounter for fracture with delayed healing
- K subsequent encounter for fracture with nonunion
- P subsequent encounter for fracture with malunion
- S sequela

Dislocation of tarsometatarsal joint

- A initial encounter
- D subsequent encounter
- S sequela

Thank You!!



Forefoot and Midfoot ICD-10 Coding

Jeffrey D. Lehrman, DPM, FASPS, MAPWCA, CPC

Advisor, APMA Coding Committee

Advisor, APMA MACRA Task Force

Expert Panelist, Codingline

Fellow, American Academy of Podiatric Practice Management

Board of Directors, American Society of Podiatric Surgeons

Board of Directors, American Professional Wound Care Association

Editorial Advisory Board, *WOUNDS*

Twitter: [@DrLehrman](https://twitter.com/DrLehrman)